



Article

Human Breathable Air in a Mediterranean Forest: Characterization of Monoterpene Concentrations under the Canopy

Albert Bach ^{1,2,*} , Ana Maria Yáñez-Serrano ^{3,4} , Joan Llusia ^{3,4}, Iolanda Filella ^{3,4}, Roser Maneja ^{2,5,6} and Josep Penuelas ^{3,4}

- ¹ Institute of Environmental Science and Technology (ICTA), Autonomous University of Barcelona (UAB), Z Building, ICTA-ICP, Carrer de les columnes, UAB Campus, Bellaterra (Cerdanyola del Vallès), 08193 Barcelona, Spain
 - ² Environment and Human Health Laboratory (EH² Lab), Forest Science and Technology Center of Catalonia, Crta. de St. Llorenç de Morunys, km 2, 25280 Solsona, Spain; roser.maneja@uab.cat
 - ³ CREAF, Campus Universitat Autònoma de Barcelona, Cerdanyola del Vallès, 08193 Barcelona, Catalonia, Spain; a.yanez@creaf.uab.cat (A.M.Y.-S.); j.llusia@creaf.uab.cat (J.L.); iola@creaf.uab.cat (I.F.); josep.penuelas@uab.cat (J.P.)
 - ⁴ CSIC, Global Ecology Unit CREAF-CSIC-UAB, Bellaterra, 08193 Barcelona, Catalonia, Spain
 - ⁵ Forest Science and Technology Center of Catalonia, Crta. de St. Llorenç de Morunys, km 2, 25280 Solsona, Spain
 - ⁶ Geography Department, Autonomous University of Barcelona (UAB), B Building, UAB Campus, Bellaterra (Cerdanyola del Vallès), 08193 Barcelona, Spain
- * Correspondence: albert.bach@uab.cat; Tel.: +34-935868654

Received: 14 May 2020; Accepted: 16 June 2020; Published: 18 June 2020



Abstract: Monoterpenes have been identified as potential determinants of the human health effects induced by forest exposure. The present study characterizes the total monoterpene concentrations at nose height in a Mediterranean Holm oak forest located in North-East Iberian Peninsula during the annual emission peak (summer and autumn: June to November) using a Proton Transfer Reaction–Mass Spectrometry (PTR-MS). Results show a strong variability of the total monoterpene concentrations in season and daytime. The concentration peak appears during July and August. These two months displayed two average maxima in their diel cycles: One during early morning (from 6:00 to 8:00, 0.30 ppbv for July and 0.41 ppbv for August) and another one at early afternoon (from 13:00 to 15:00, 0.27 ppbv during July and 0.32 ppbv during August). Monoterpene concentrations were strongly related with the temperature (exponentially) and solar radiation (rectangular hyperbolic relationship). The concentrations registered here are similar or higher than in previous ex situ studies showcasing the effects of forests on human health. These findings provide relevant data for the scientific and healthcare community by improving the understanding of monoterpene dynamics at nose height and suggesting further research on the effects of forests on human health, particularly in the Mediterranean region.

Keywords: monoterpenes; forest exposure; PTR-MS; forest chemistry; BVOCs; forest bathing

1. Introduction

Interest in understanding if and how the exposure to forests leads to human health benefits is growing among the studies analyzing the effects of nature on human health [1–4]. However, evidence connecting particular forest variables and health effects is still inconclusive, due to the lack of forest descriptions and the high heterogeneity of approaches and results within the studies performed

so far [5]. Furthermore, Mediterranean forests appear to be scarcely studied in comparison with Asian ones [5], where abiotic variables and forest ecosystems may differ. This growing body of research has argued that biogenic volatile organic compounds (BVOCs) may partly explain the health effects induced by forest exposure [6–9]. BVOCs emitted by forests have diverse roles at multiple scales; from cellular protection and defense at the foliar level, through to chemical signaling at the regional level, up to influencing rainfall at the ecosystems scale [10]. These compounds profoundly affect biosphere–atmosphere interactions by atmospheric reactivity, aerosol growth processes, cloud formation, and therefore radiative balance [11,12].

Monoterpenes are part of the isoprenoid class, the largest class of BVOCs and major components of the forest atmospheres [13]. These compounds are produced by plants as a defensive mechanism against environmental stress and herbivory [13–15]. Additionally, monoterpenes are highly reactive compounds that have a high yield in secondary organic aerosol formation [16]. Monoterpene emission vary across plant species and functional type [17,18], which further changes with an altered physiological plant state [19], leading to distinct spatial [20] and temporal [21,22] emissions worldwide. Monoterpenes have been shown to interact with human health in different ways, from stress relief to an effect on the immune function. Inhaling monoterpenes has been shown to decrease blood pressure and cortisol levels [23], improve antibiotics efficiency [24], or boost the immune system [8,25,26], especially by increasing the percentage and activity of natural killer cells [7,9,27,28]. Furthermore, monoterpenes that relevantly affect cellular and animal systems have shown anti-inflammatory, antitumorigenic, or neuroprotective activities [29].

To our knowledge, only one study has analyzed the absorption of monoterpenes in blood after forest exposure, by identifying the monoterpene species present in a coniferous forest atmosphere in serum samples of individuals after walking in the forest [30]. The study demonstrated an increase in the amount of α -pinene in the serum after the individuals walked in the forest as well as differences in monoterpene composition and abundance between coniferous and broad-leaved forest air [30]. In addition, Lee et al. (2018) reported that the mean atmospheric concentration of monoterpenes at a 1.5-m height was higher in a natural than a tended forest [8], suggesting that vegetation characteristics derived from management may affect terpenes' concentrations under the canopy.

While studies are increasingly providing evidence on the role of monoterpenes in human health, research is needed to describe and predict the composition and abundance of monoterpenes at nose height under forest canopies, analyze the absorption and metabolism of these chemicals by humans, and identify the mechanisms leading to health effects. To address this gap, this article characterized the total monoterpene concentrations at nose height in a Mediterranean holm oak forest during the annual emission peak and posterior months and sheds light on the potential effects of such concentrations on human health.

2. Materials and Methods

2.1. Measurement Site

The study site was located in Montseny Natural Park—Biosphere Reserve (NE Iberian Peninsula, Spain) in a Holm oak forest dominated by *Quercus ilex* (1100 per ha), which is highly representative of the montane holm oak (*Quercus ilex* L.) forests in the northern Mediterranean regions [31], Figure 1. At Montseny, due to coppicing until the 1950s and the posterior selective thinning, holm oaks have nowadays become dense forest of resprout origin [32]. The average diameter at breast height (DBH) for *Quercus ilex* was 16.36 ± 0.917 cm. Other species like *Pinus halepensis* or *Erica arborea* were identified in the study area but not abundantly. Concerning the forest structure, vegetation cover at the 8-m height was 100% while the layers below did not exceed coverages of 10% except for heights of 0.25 and 0.5 m, where the understory was dense and presented coverages of 80%. This forest area has an altitude range from 415 to 550 m.




Figure 1. Sampling site.

The habitats within a 500-m radius around our sampling site are presented in Table 1 and illustrated in Figure 2.

Table 1. Habitat type, surface, and percentages of the total surface per habitat within a 500-m radius from the sampling point.

Habitat	Surface (ha)	Percentage (%)
Holm Oak Forest	33.63	43.50
Cropland	16.83	21.77
Mixed forest (holm oak + deciduous oaks)	9.97	12.90
Urban and industrial areas	5.41	6.99
Bushland	4.94	6.39
Riparian vegetation	4.19	5.42
Cork oak forest	2.34	3.03
TOTAL	77.31	




Figure 2. Study area: Sampling point, buffers (500 m, 1 and 2 km), and habitats within the zone.

2.2. Meteorological Conditions

At the sampling site (41°43'43.29" N, 2°26'24.35" E, 422 m a.s.l.), there was a manual meteorological station with daily values of temperature at 8:00 am, and daily maximum and minimum temperature and precipitation. In order to complement the data with higher frequency datasets, we obtained meteorological data from two other stations of the Servei Meteorològic de Catalunya (i.e., Catalonia's Meteorological Service): Radiation and humidity from Puig Sesolles station (41°46'25.00" N, 2°26'15.90" E, 1668 m a.s.l.) 5 km away from our sampling site, and temperature from Tagamanent station (41°44'51.43" N, 2°18'10.64" E, 1030 m a.s.l.) 12 km away from the sampling site. The vapor pressure deficit was calculated with the solar radiation data from Puig Sesolles and the temperature data from Tagamanent.

2.3. PTR-MS Sampling Methodology

Measurements of the total monoterpene concentrations were performed with a PTR-MS (Ionicon Analytic GmbH, Innsbruck, Austria) from 26 June to 15 November 2019. One isolated 10-m Teflon tube (OD $\frac{1}{4}$) was installed at 1.5 m attached with a rope to a holm oak tree. Air was sucked to inside the Park Office where the PTR-MS was located by a pump (JUN-AIR, Benton Harbor, MI, USA). The PTR-MS was operated at standard conditions (2.2 mbar drift pressure, 600 V drift voltage, 127 Td) [33]. A catalytic converter (Supelco Inc., Bellefonte, USA with platinum pellets heated to 380 °C) was used to monitor the background impurities of the PTR-MS. The background signal for each compound was calculated once every hour. Background values were interpolated over the time of the measurements. Humidity-dependent calibrations (using bubbled zero air to dilute the standard, regulated as close as possible to ambient humidity conditions) were performed using a gravimetrically prepared multicomponent standard, including methanol (m/z 33), acetonitrile (m/z 42), acetaldehyde (m/z 45), acetone (m/z 59), isoprene (m/z 69), MACR (m/z 71), MEK (m/z 73), and α -pinene (m/z 137), with several dilution steps (Riemer Environmental Inc., Miami, USA).

The PTR-MS technique separates per mass and not per compound, so other compounds than monoterpenes with the same mass might be measured together. However, sporadic GC-MS screening at the site confirmed the presence of monoterpenes (i.e., m/z 137) in ambient air, more specifically α -pinene and β -pinene.

2.4. Data Analysis

Igor Pro (Wavemetrics Inc., Portland, OR, USA) was used for the calculations of BVOCs concentrations, data time series treatment, graphing, and descriptive statistics. We conducted a generalized linear model (GLM) to determine the effect of the available atmospheric data (temperature, precipitation, relative humidity, solar radiation, ozone, and vapor pressure deficit) on monoterpene concentrations using R (Version 1.2.5033). Monoterpene concentrations were log-transformed to ensure normality of the residuals. A $p < 0.05$ was considered significant.

3. Results

Figure 3 shows the complete time series of the total monoterpene concentrations measured in this study from June to November 2019, which ranged from 0.02 to 0.13 ppbv (25 percentile and 75 percentile), with maximum values reaching 1.24 ppbv. Total monoterpenes had a clear seasonality with higher concentrations during summer as well as a strong daily pattern with highest concentrations during daylight.




Figure 3. Total monoterpene air concentrations during the sampling period (from June to November 2019). The gaps are due to instrument lack of power in the sampling site.

The average monthly diel cycles (Figure 4) showed that August and July had the highest concentrations. These two months displayed two average maxima in their diel cycles, one during early morning (from 6:00 to 8:00, 0.30 ppbv for July and 0.41 ppbv for August) and another one at early afternoon (from 13:00 to 15:00, 0.27 ppbv during July and 0.32 ppbv during August). These peaks were not identified in the rest of the sampled months (September, October, and November), where total monoterpene concentrations increased with the solar radiation and a peak around 14:00, coinciding with the temperature diel cycle peak. Additionally, concentrations values stayed relatively constant during night, with 0.05 ppbv for July and August decreasing to 0.02 ppbv in November.




Figure 4. Half-hour average diel cycle of total monoterpene concentrations per month. Error bar represents the standard error.

Additionally, Figure 5 shows the monoterpene concentrations were strongly related with temperature (exponentially) and solar radiation (rectangular hyperbolic relationship). The evolution of the meteorological variables correlated with the monoterpenes is presented in Figure S1. The GLM analysis demonstrated a significant effect of solar radiation (t -value = 22.770; p -value = 2×10^{-16}),

temperature (t -value = 21.645; p -value = 2×10^{-16}), humidity (t -value = 9.506; p -value = 2×10^{-16}), and vapor pressure deficit (vpd) (t -value = 6.236; p -value = 4.82×10^{-16}) (Figures S2 and S3 and Table S1).




Figure 5. 3-D relationship of monoterpene concentrations with temperature and radiation.

4. Discussion

The present study analyzed the under-canopy variations of monoterpene concentrations in a Mediterranean holm oak forest in Montseny Natural Park—Biosphere Reserve (NE Iberian Peninsula) continuously from June to November 2019 using a PTR-MS. This is the longest time series of total monoterpene concentrations ever measured in a Mediterranean holm oak forest at the nose level. During the sampled period, we identified seasonal and daily patterns. The highest concentrations occurred during summer, and diel cycles changed among months, with peaks in early morning in summer and peaks at midday in the entire period. The total monoterpene variations were significantly linked to abiotic/atmospheric variables, such as temperature and solar radiation, as well as to humidity and vapor pressure deficit (vpd).

The highest monoterpene concentrations occurred during July and August. These results are consistent with other studies that found higher monoterpene concentrations above the canopy level in summer in an holm oak forest (*Quercus ilex* L.) from the same Mediterranean region [34,35]. Previous studies in other regions have as well observed higher monoterpene emissions during summer period [36,37]. Furthermore, monoterpene emission rates measured at the leaf or branch level have proved to be higher from July to August in *Quercus ilex* [34,35,38,39]. This seasonality of monoterpene concentrations is associated to the seasonality of plant monoterpene emission [19,40] mostly linked to the seasonal changes in temperature [41,42] and other meteorological variables, such as solar radiation [17,43,44], which explains as well the decrease during the autumn months (October and November).

We found higher monoterpene concentrations during daytime, with maxima at early morning (from 6:00 to 8:00) and at early afternoon (from 13:00 to 15:00). This early morning peak has been previously reported at a nearby site above forest canopies [38,45,46]. This peak has been attributed to the biogenic monoterpene emissions at the site, which have not yet been influenced by mountain

breeze (which starts a few hours later in the day) [46]. The peak at around 14:00 is also related to the biogenic origin of monoterpenes, which is enhanced at maximum temperature and solar radiation. In fact, the maximum at 14:00 coinciding with the temperature peak has been reported previously for areas with light- and temperature-dependent monoterpene-emitting trees, such as tropical trees [47,48]. This differs from other forest sites, including Mediterranean, that documented the diel cycle of the monoterpene concentration to peak at nighttime [37,49,50]. This inconsistency is due to the fact that holm oaks emit monoterpenes in a light- and temperature-dependent manner (so called *de novo* biosynthesis) and not from storage pools at night, like boreal coniferous species [17,43], and thus maximum emissions occur during the middle of the day [51].

Monoterpene emissions are well-known to be affected by atmospheric variables, such as air temperature, solar radiation, and air humidity [17,43,44]. Our results from the GLM analysis support this fact by showing a strong effect of temperature, solar radiation, humidity, and vpd on monoterpene concentrations. Ambient concentrations of a volatile organic compound can be affected by the strength of emissions (either biogenic or anthropogenic), the effectiveness of air mixing, and the strength of sink processes, including deposition and chemical degradation. However, the strong effect of temperature and solar radiation on the observed concentrations suggests the dominance of a biogenic source for total monoterpenes. Further on, as the site is dominated by *Quercus ilex* forests, a dominant monoterpene emitter tree species, we assume that the main origin for the observed concentrations comes from the monoterpene emission of this tree species. This further supports the biogenic origin of the measured concentrations, similar to a previous study at a nearby site [46]. Furthermore, the daily patterns of the concentrations could as well be partly associated to atmospheric vertical stability and wind speed and direction as shown in previous studies [52]. Reasonable interpolation from existing weather stations in this regard could not be provided in the present study, but further research is warranted at this regard.

Our results report similar monoterpene concentrations to another study that has measured these compounds under coniferous forest canopies in Japan [53]. The most abundant monoterpenes in Mediterranean holm oak forests are α -pinene, sabinene, limonene, camphene, and α -phellandrene [38]. Some of these monoterpene species have been previously found to be absorbed by humans after a 60-min walk through a coniferous forest with a six-fold increase of the α -pinene peak in blood after forest exposure [30]. These monoterpenes have been related with physiological changes in humans, in particular α -phellandrene has been significantly associated with an acute insulin reaction [8]. Some *ex situ* and *in vitro* studies have shown enhancing effects of monoterpenes on the human immune system but at higher monoterpene concentrations than those we registered in our study (approximately from 50 to 1900 ppbv) [7,9]. Nevertheless, the concentrations we report here (from 114 to 725 ngm⁻³ during the whole sampling period and up to 2230 ngm⁻³ in August) happen to be comparable [53] or much greater [27,28] than in previous *in situ* studies that identified similar health outcomes to the ones observed in the *in vitro* and *ex situ* experiments. The concentrations of monoterpenes of our studied Mediterranean forest are also higher than other *in situ* studies characterizing the forest atmosphere under the canopy [54] or testing the effects of forests on other human systems and functions [8]. Taken together, these findings suggest that the Mediterranean holm oak forests constitute a suitable forest environment to develop further research on the effects of monoterpenes on human health.

5. Conclusions

This study analyzed the monoterpene concentrations at nose height in a Mediterranean Holm oak forest from June to November. We identified a strong variability of the total monoterpene concentrations in season and daytime with its peak during summer. Additionally, the monthly average diel cycles showed two main peaks at early morning (from 6:00 to 8:00) and early afternoon (from 13:00 to 18:00) during July and August. Monoterpene concentrations significantly correlated with air temperature, solar radiation, air humidity, and vapor pressure deficit (vpd).

If BVOCS play a key role in the interaction between forest and human health as highlighted in the literature, our analysis provided relevant information concerning the availability of these compounds

at the nose level, thus suggesting that humans walking in the studied forest may be subject to a potentially high absorption of monoterpenes into their bloodstream, especially at early morning and from midday. Our findings contribute to improving the understanding of monoterpene dynamics under the canopy and enhancing the development of more applied research on the effects of forests on human health, particularly in the Mediterranean region.

Supplementary Materials: The following are available online at <http://www.mdpi.com/1660-4601/17/12/4391/s1>, Figure S1: Evolution of the meteorological variables during the sampling period; Figure S2: Correlation between air humidity and monoterpenes concentrations; Figure S3: Correlation between air vapor pressure deficit (vpd) and monoterpenes concentrations; and Table S1: Outcomes from the GLM conducted to assess the effect of air temperature, solar radiation, air humidity and vapor pressure deficit (vpd) on monoterpenes concentrations.

Author Contributions: Conceptualization, A.M.Y.-S., A.B. and J.P.; methodology, A.M.Y.-S. and A.B.; software, A.M.Y.-S.; validation, I.F. and J.L.; formal analysis, A.M.Y.-S. and A.B.; data curation, J.P. and I.F.; writing—original draft preparation, A.M.Y.-S. and A.B.; writing—review and editing, A.M.Y.-S., A.B., R.M., J.L., I.F. and J.P.; supervision, R.M. and J.P. All authors have read and agreed to the published version of the manuscript.

Funding: The research leading to these results has received funding from “la Caixa” Foundation. Furthermore, this study was supported by the Spanish Government Grant CGL2013-48074-P, the Catalan Government grant SGR 2017-1005, the European Research Council Synergy grant ERC-2013-SyG 610028 IMBALANCE-P.

Acknowledgments: We truly thank David Bartolomeu for his field assistance during the measurements. We would like to especially thank Joana Barber and Daniel Guinart from the Natural Park—Biosphere Reserve of Montseny for allowing us to use the Park facilities for the study. We are especially grateful for Esteve Corbera’s relevant contributions to the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Tsunetsugu, Y.; Park, B.J.; Miyazaki, Y. Trends in research related to “shinrin-yoku” (taking in the forest atmosphere or forest bathing) in Japan. *Environ. Health Prev. Med.* **2010**, *15*, 27–37. [[CrossRef](#)] [[PubMed](#)]
2. Lee, J.; Park, B.J.; Tsunetsugu, Y.; Ohira, T.; Kagawa, T.; Miyazaki, Y. Effect of forest bathing on physiological and psychological responses in young Japanese male subjects. *Public Health* **2011**, *125*, 93–100. [[CrossRef](#)] [[PubMed](#)]
3. Song, C.; Ikei, H.; Miyazaki, Y. Physiological effects of nature therapy: A review of the research in Japan. *Int. J. Environ. Res. Public Health* **2016**, *13*, 781. [[CrossRef](#)] [[PubMed](#)]
4. Park, B.J.; Tsunetsugu, Y.; Kasetani, T.; Kagawa, T.; Miyazaki, Y. The physiological effects of Shinrin-yoku (taking in the forest atmosphere or forest bathing): Evidence from field experiments in 24 forests across Japan. *Environ. Health Prev. Med.* **2010**, *15*, 18–26. [[CrossRef](#)] [[PubMed](#)]
5. Bach, A.; Peñuelas, J.; Clarà, J.; Llusia, J.; Campillo, I.; López, F.; Maneja, R. How Should Forests Be Characterized in Regard to Human Health? Evidence from Existing Literature. *Int. J. Environ. Res. Public Health* **2020**, *17*, 1027. [[CrossRef](#)]
6. Nazaruk, J.; Borzym-Kluczyk, M. The role of triterpenes in the management of diabetes mellitus and its complications. *Phytochem. Rev.* **2015**, *14*, 675–690. [[CrossRef](#)]
7. Li, Q.; Kobayashi, M.; Wakayama, Y.; Inagaki, H.; Katsumata, M.; Hirata, Y.; Hirata, K.; Shimizu, T.; Kawada, T.; Park, B.J.; et al. Effect of phytoncide from trees on human natural killer cell function. *Int. J. Immunopathol. Pharmacol.* **2009**, *22*, 951–959. [[CrossRef](#)]
8. Lee, K.J.; Hur, J.; Yang, K.S.; Lee, M.K.; Lee, S.J. Acute Biophysical Responses and Psychological Effects of Different Types of Forests in Patients With Metabolic Syndrome. *Environ. Behav.* **2018**, *50*, 298–323. [[CrossRef](#)]
9. Li, Q.; Nakadai, A.; Matsushima, H.; Miyazaki, Y.; Krensky, A.; Kawada, T.; Morimoto, K. Phytoncides (wood essential oils) induce human natural killer cell activity. *Immunopharmacol. Immunotoxicol.* **2006**, *28*, 319–333. [[CrossRef](#)]
10. Laothawornkitkul, J.; Taylor, J.E.; Paul, N.D.; Hewitt, C.N. Biogenic volatile organic compounds in the Earth system. *New Phytol.* **2009**, *183*, 27–51. [[CrossRef](#)]
11. Kulmala, M.; Nieminen, T.; Chellapermal, R.; Makkonen, R.; Bäck, J.; Kerminen, V.-M. Climate Feedbacks Linking the Increasing Atmospheric CO₂ Concentration, BVOC Emissions, Aerosols and Clouds in Forest Ecosystems. In *Biology, Controls and Models of Tree Volatile Organic Compound Emissions*; Springer: Dordrecht, The Netherlands, 2013; pp. 489–508.

12. Pöschl, U.; Martin, S.T.; Sinha, B.; Chen, Q.; Gunthe, S.S.; Huffman, J.A.; Borrmann, S.; Farmer, D.K.; Garland, R.M.; Helas, G.; et al. Rainforest Aerosols as Biogenic Nuclei of Clouds and Precipitation in the Amazon. *Science* **2010**, *329*, 1513–1516. [[CrossRef](#)] [[PubMed](#)]
13. Gershenzon, J.; Dudareva, N. The function of terpene natural products in the natural world. *Nat. Chem. Biol.* **2007**, *3*, 408–414. [[CrossRef](#)] [[PubMed](#)]
14. Mewalal, R.; Rai, D.K.; Kainer, D.; Chen, F.; Külheim, C.; Peter, G.F.; Tuskan, G.A. Plant-Derived Terpenes: A Feedstock for Specialty Biofuels. *Trends Biotechnol.* **2017**, *35*, 227–240. [[CrossRef](#)] [[PubMed](#)]
15. Martin, D.M.; Gershenzon, J.; Bohlmann, J. Induction of volatile terpene biosynthesis and diurnal emission by methyl jasmonate in foliage of Norway spruce. *Plant Physiol.* **2003**, *132*, 1586–1599. [[CrossRef](#)] [[PubMed](#)]
16. Zhao, D.F.; Kaminski, M.; Schlag, P.; Fuchs, H.; Acir, I.-H.; Bohn, B.; Häsel, R.; Kiendler-Scharr, A.; Rohrer, F.; Tillmann, R.; et al. Secondary organic aerosol formation from hydroxyl radical oxidation and ozonolysis of monoterpenes. *Atmos. Chem. Phys.* **2015**, *15*, 991–1012. [[CrossRef](#)]
17. Kesselmeier, J.; Staudt, M. Biogenic volatile organic compounds (VOC): An overview on emission, physiology and ecology. *Environ. Pollut.* **2000**, *109*, 175.
18. Guenther, A.B.; Jiang, X.; Heald, C.L.; Sakulyanontvittaya, T.; Duhl, T.; Emmons, L.K.; Wang, X. The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): An extended and updated framework for modeling biogenic emissions. *Geosci. Model Dev.* **2012**, *5*, 1471–1492. [[CrossRef](#)]
19. Harrison, S.P.; Morfopoulos, C.; Dani, K.G.S.; Prentice, I.C.; Arneth, A.; Atwell, B.J.; Barkley, M.P.; Leishman, M.R.; Loreto, F.; Medlyn, B.E.; et al. Volatile isoprenoid emissions from plastid to planet. *New Phytol.* **2013**, *197*, 49–57. [[CrossRef](#)] [[PubMed](#)]
20. Sindelarova, K.; Granier, C.; Bouarar, I.; Guenther, A.; Tilmes, S.; Stavrou, T.; Müller, J.-F.; Kuhn, U.; Stefani, P.; Knorr, W. Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years. *Atmos. Chem. Phys.* **2014**, *14*, 9317–9341. [[CrossRef](#)]
21. Hantson, S.; Knorr, W.; Schurgers, G.; Pugh, T.A.M.; Arneth, A. Global isoprene and monoterpene emissions under changing climate, vegetation, CO₂ and land use. *Atmos. Environ.* **2017**, *155*, 35–45. [[CrossRef](#)]
22. Peñuelas, J.; Staudt, M. BVOCs and global change. *Trends Plant Sci.* **2010**, *15*, 133–144. [[CrossRef](#)] [[PubMed](#)]
23. Nam, E.-S.; Uhm, D.-C. Effects of Phytoncides Inhalation on Serum Cortisol Level and Life Stress of College Students. *Korean J. Adult Nurs.* **2008**, *20*, 697–706.
24. Sieniawska, E.; Sawicki, R.; Swatko-Ossor, M.; Napiorkowska, A.; Przekora, A.; Ginalska, G.; Swatko-Ossor, M.; Augustynowicz-Kopec, E. The Effect of Combining Natural Terpenes and Antituberculous Agents against Reference and Clinical Mycobacterium tuberculosis Strains. *Molecules* **2018**, *23*, 176. [[CrossRef](#)] [[PubMed](#)]
25. Komori, T.; Fujiwara, R.; Tanida, M.; Nomura, J.; Yokoyama, M.M. Effects of citrus fragrance on immune function and depressive states. *Neuroimmunomodulation* **1995**, *2*, 174–180. [[CrossRef](#)]
26. Lin, J.J.; Lu, K.W.; Ma, Y.S.; Tang, N.Y.; Wu, P.P.; Wu, C.C.; Lu, H.F.; Lin, J.G.; Chung, J.G. Alpha-phellandrene, a natural active monoterpene, influences a murine WEHI-3 leukemia model in vivo by enhancing macrophage phagocytosis and natural killer cell activity. *In Vivo (Brooklyn)* **2014**, *28*, 583–588.
27. Li, Q.; Morimoto, K.; Nakadai, A.; Inagaki, H.; Katsumata, M.; Shimizu, T.; Hirata, Y.; Hirata, K.; Suzuki, H.; Miyazaki, Y.; et al. Forest bathing enhances human natural killer activity and expression of anti-cancer proteins. *Int. J. Immunopathol. Pharmacol.* **2007**, *20*, 3–8. [[CrossRef](#)] [[PubMed](#)]
28. Li, Q.; Morimoto, K.; Kobayashi, M.; Inagaki, H.; Katsumata, M.; Hirata, Y.; Hirata, K.; Shimizu, T.; Li, Y.J.; Wakayama, T.; et al. A forest bathing trip increases human natural killer activity and expression of anti-cancer proteins in female subjects. *J. Biol. Regul. Homeost. Agents* **2008**, *22*, 45–55. [[PubMed](#)]
29. Cho, K.S.; Lim, Y.R.; Lee, K.; Lee, J.; Lee, J.H.; Lee, I.S. Terpenes from forests and human health. *Toxicol. Res.* **2017**, *33*, 97–106. [[CrossRef](#)] [[PubMed](#)]
30. Sumitomo, K.; Akutsu, H.; Fukuyama, S.; Minoshima, A.; Kukita, S.; Yamamura, Y.; Sato, Y.; Hayasaka, T.; Osanai, S.; Funakoshi, H.; et al. Conifer-Derived Monoterpenes and Forest Walking. *Mass Spectrom. (Tokyo, Japan)* **2015**, *4*, A0042. [[CrossRef](#)]
31. Terradas, J. Holm Oak and Holm Oak Forests: An Introduction. In *Ecology of Mediterranean Evergreen oak Forests*; Springer: Berlin/Heidelberg, Germany, 1999; pp. 3–14.
32. Avila, A.; Rodrigo, A. Trace metal fluxes in bulk deposition, throughfall and stemflow at two evergreen oak stands in NE Spain subject to different exposure to the industrial environment. *Atmos. Environ.* **2004**, *38*, 171–180. [[CrossRef](#)]

33. Lindinger, W.; Jordan, A. Proton-transfer-reaction mass spectrometry (PTR-MS): On-line monitoring of volatile organic compounds at pptv levels. *Chem. Soc. Rev.* **1998**, *27*, 347. [[CrossRef](#)]
34. Peñuelas, J.; Guenther, A.; Rapparini, F.; Llusia, J.; Filella, I.; Seco, R.; Estiarte, M.; Mejia-Chang, M.; Ogaya, R.; Ibañez, J.; et al. Intensive measurements of gas, water, and energy exchange between vegetation and troposphere during the MONTES campaign in a vegetation gradient from short semi-desertic shrublands to tall wet temperate forests in the NW Mediterranean Basin. *Atmos. Environ.* **2013**, *75*, 348–364. [[CrossRef](#)]
35. Seco, R.; Peñuelas, J.; Filella, I.; Llusia, J.; Molowny-Horas, R.; Schallhart, S.; Metzger, A.; Müller, M.; Hansel, A. Contrasting winter and summer VOC mixing ratios at a forest site in the Western Mediterranean Basin: The effect of local biogenic emissions. *Atmos. Chem. Phys.* **2011**, *11*, 13161–13179. [[CrossRef](#)]
36. Ruuskanen, T.M.; Taipale, R.; Rinne, J.; Kajos, M.K.; Hakola, H.; Kulmala, M. Quantitative long-term measurements of VOC concentrations by PTR-MS: Annual cycle at a boreal forest site. *Atmos. Chem. Phys. Discuss.* **2009**, *9*, 81–134. [[CrossRef](#)]
37. Jordan, C.; Fitz, E.; Hagan, T.; Sive, B.; Frinak, E.; Haase, K.; Cottrell, L.; Buckley, S.; Talbot, R. Long-term study of VOCs measured with PTR-MS at a rural site in New Hampshire with urban influences. *Atmos. Chem. Phys.* **2009**, *9*, 4677–4697. [[CrossRef](#)]
38. Llusia, J.; Peñuelas, J.; Seco, R.; Filella, I. Seasonal changes in the daily emission rates of terpenes by *Quercus ilex* and the atmospheric concentrations of terpenes in the natural park of Montseny, NE Spain. *J. Atmos. Chem.* **2012**, *69*, 215–230. [[CrossRef](#)]
39. Llusia, J.; Peñuelas, J.; Guenther, A.; Rapparini, F. Seasonal variations in terpene emission factors of dominant species in four ecosystems in NE Spain. *Atmos. Environ.* **2013**, *70*, 149–158. [[CrossRef](#)]
40. Peñuelas, J.; Llusia, J. The complexity of factors driving volatile organic compound emissions by plants. *Biol. Plant.* **2001**, *44*, 481–487. [[CrossRef](#)]
41. Loreto, F.; Forster, A.; Durr, M.; Csiky, O.; Seufert, G. On the monoterpene emission under heat stress and on the increased thermotolerance of leaves of *Quercus ilex* L. fumigated with selected monoterpenes. *Plant Cell Environ.* **1998**, *21*, 101–107. [[CrossRef](#)]
42. Loreto, F.; Pinelli, P.; Manes, F.; Kollist, H. Impact of ozone on monoterpene emissions and evidence for an isoprene-like antioxidant action of monoterpenes emitted by *Quercus ilex* leaves. *Tree Physiol.* **2004**, *24*, 361–367. [[CrossRef](#)]
43. Llusia, J.; Peñuelas, J. *Pinus halepensis* and *Quercus ilex* terpene emission as affected by temperature and humidity. *Biol. Plant.* **1999**, *42*, 317–320. [[CrossRef](#)]
44. Staudt, M.; Rambal, S.; Joffre, R.; Kesselmeier, J. Impact of drought on seasonal monoterpene emissions from *Quercus ilex* in southern France. *J. Geophys. Res. Atmos.* **2002**, *107*, ACH 15-1–ACH 15-9. [[CrossRef](#)]
45. Seco, R.; Filella, I.; Llusia, J.; Peñuelas, J. Methanol as a signal triggering isoprenoid emissions and photosynthetic performance in *Quercus ilex*. *Acta Physiol. Plant.* **2011**, *33*, 2413–2422. [[CrossRef](#)]
46. Seco, R.; Peñuelas, J.; Filella, I.; Llusia, J.; Schallhart, S.; Metzger, A.; Müller, M.; Hansel, A. Volatile organic compounds in the western Mediterranean basin: Urban and rural winter measurements during the DAURE campaign. *Atmos. Chem. Phys.* **2013**, *13*, 4291–4306. [[CrossRef](#)]
47. Yáñez-Serrano, A.M.; Nölscher, A.C.; Williams, J.; Wolff, S.; Alves, E.G.; Martins, G.A.; Bourtsoukidis, E.; Brito, J.; Jardine, K.; Artaxo, P.; et al. Diel and seasonal changes of biogenic volatile organic compounds within and above an Amazonian rainforest. *Atmos. Chem. Phys.* **2015**, *15*, 3359–3378. [[CrossRef](#)]
48. Jardine, A.B.; Jardine, K.J.; Fuentes, J.D.; Martin, S.T.; Martins, G.; Durgante, F.; Carneiro, V.; Higuchi, N.; Manzi, A.O.; Chambers, J.Q. Highly reactive light-dependent monoterpenes in the Amazon. *Geophys. Res. Lett.* **2015**, *42*, 1576–1583. [[CrossRef](#)]
49. Davison, B.; Taipale, R.; Langford, B.; Misztal, P.; Fares, S.; Matteucci, G.; Loreto, F.; Cape, J.N.; Rinne, J.; Hewitt, C.N. Concentrations and fluxes of biogenic volatile organic compounds above a Mediterranean macchia ecosystem in western Italy. *Biogeosciences* **2009**, *6*, 1655–1670. [[CrossRef](#)]
50. Yassaa, N.; Song, W.; Lelieveld, J.; Vanhatalo, A.; Bäck, J.; Williams, J. Diel cycles of isoprenoids in the emissions of Norway spruce, four Scots pine chemotypes, and in Boreal forest ambient air during HUMPPA-COPEC-2010. *Atmos. Chem. Phys.* **2012**, *12*, 7215–7229. [[CrossRef](#)]
51. Staudt, M.; Bertin, N. Light and temperature dependence of the emission of cyclic and acyclic monoterpenes from holm oak (*Quercus ilex* L.) leaves. *Plant Cell Environ.* **1998**, *21*, 385–395. [[CrossRef](#)]
52. Meneguzzo, F.; Albanese, L.; Bartolini, G.; Zabini, F. Temporal and spatial variability of volatile organic compounds in the forest atmosphere. *Int. J. Environ. Res. Public Health* **2019**, *16*, 4915. [[CrossRef](#)] [[PubMed](#)]

53. Li, Q.; Morimoto, K.I.; Kobayashi, M.; Inagaki, H.; Katsumata, M.; Hirata, Y.; Hirata, K.; Suzuki, H.; Li, Y.; Wakayama, Y.; et al. Visiting a forest, but not a city, increases human natural killer activity and expression of anti-cancer proteins. *Int. J. Immunopathol. Pharmacol.* **2008**, *21*, 117–127. [[CrossRef](#)] [[PubMed](#)]
54. Kim, G.; Park, B.J.; Yeon, P.S.; Lee, S.; Joung, D.; Park, C.; Koga, S. Case Study on the Changes in the Physical Environment in Forest Healing Spaces. *J. Fac. Agr. Kyushu Univ.* **2016**, *61*, 375–381.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).